

Accumulating Data to Optimally Predict Obesity Treatment (ADOPT) Core Measures: Psychosocial Domain

Angelina R. Sutin¹, Kerri Boutelle², Susan M. Czajkowski³, Elissa S. Epel⁴, Paige A. Green³, Christine M. Hunter⁵, Elise L. Rice³, David M. Williams⁶, Deborah Young-Hyman⁵, and Alexander J. Rothman⁷

Background: Within the Accumulating Data to Optimally Predict obesity Treatment (ADOPT) Core Measures Project, the psychosocial domain addresses how psychosocial processes underlie the influence of obesity treatment strategies on weight loss and weight maintenance. The subgroup for the psychosocial domain identified an initial list of high-priority constructs and measures that ranged from relatively stable characteristics about the person (cognitive function, personality) to dynamic characteristics that may change over time (motivation, affect).

Objectives: This paper describes (a) how the psychosocial domain fits into the broader model of weight loss and weight maintenance as conceptualized by ADOPT; (b) the guiding principles used to select constructs and measures for recommendation; (c) the high-priority constructs recommended for inclusion; (d) domain-specific issues for advancing the science; and (e) recommendations for future research.

Significance: The inclusion of similar measures across trials will help to better identify how psychosocial factors mediate and moderate the weight loss and weight maintenance process, facilitate research into dynamic interactions with factors in the other ADOPT domains, and ultimately improve the design and delivery of effective interventions.

Obesity (2018) **26**, S45–S54. doi:10.1002/oby.22160

Introduction

Approximately two-thirds of adults in the United States are actively trying to manage their weight, either through losing weight or maintaining the same weight (1). Despite this widespread effort to achieve a healthier weight, intervention strategies have had limited success in producing long-term weight loss and maintenance (2). There is also tremendous variability in treatment response, with many treatments effective for some individuals but not others (3). Weight loss/maintenance is the result of a complex web of influences and processes that include psychosocial, biological, behavioral, and environmental factors. These pathways are often studied in isolation, but as the Accumulating Data to Optimally Predict obesity Treatment (ADOPT) Core Measures Project makes clear (4), understanding how processes within and across these domains interact is critical to identifying targets for treatment and how, when, and with whom to intervene to achieve the greatest success.

The long-term objective of the ADOPT project is to better understand the factors that contribute to individual variability in response to obesity treatment to enable more effective interventions that can be targeted and tailored to the individual. As a step toward this long-term goal, the short-term objectives are to integrate behavioral, biological, environmental, and psychosocial factors into an overarching framework for treatment response; to develop an initial list of core constructs within each domain that are critical to the process; and to identify the best measures of these constructs for recommendation for use in weight-loss trials. For the purposes of this paper, the word “construct” is used to identify mechanisms, processes, and psychological concepts broadly used and accepted in the research community and defined operationally so that they can be tested. This paper reports the recommendations of the psychosocial domain subgroup of the ADOPT working group.

¹ Department of Behavioral Sciences & Social Medicine, College of Medicine, Florida State University, Tallahassee, Florida, USA. Correspondence: Angelina R. Sutin (angelina.sutin@med.fsu.edu) ² Department of Pediatrics, University of California San Diego, San Diego, California, USA ³ National Cancer Institute, National Institutes of Health, Rockville, Maryland, USA ⁴ Department of Psychiatry, School of Medicine, University of California San Francisco, San Francisco, California, USA ⁵ Office of Behavioral and Social Sciences Research, National Institutes of Health, Bethesda, Maryland, USA ⁶ Center for Health Equity Research, School of Public Health, Brown University, Providence, Rhode Island, USA ⁷ Department of Psychology, University of Minnesota, Minneapolis, Minnesota, USA.

Funding agencies: The ADOPT Core Measures Working Group was supported by intramural funding from the following National Institutes of Health: The National Heart, Lung, and Blood Institute, the National Cancer Institute, and the Office of Disease Prevention. The Grid-Enabled Measures Database (GEM) is supported and administered by the National Cancer Institute.

Disclosure: KB has a grant from the Egg Nutrition Board. All other authors declared no conflicts of interest that are directly relevant to the work under consideration. The views expressed in this paper are those of the authors and do not necessarily represent the positions of the NIH, the DHHS, or the Federal Government.

Received: 21 December 2017; **Accepted:** 12 February 2018; **Published online** 23 March 2018. doi:10.1002/oby.22160

It has long been recognized that psychosocial factors play a role in the effectiveness of obesity treatment. The ADOPT effort seeks to capitalize on this knowledge and address previous limitations, including the need to target similar constructs across studies; use consistent, well-validated measures; and examine these constructs in conjunction with constructs from other domains. In the following sections, we describe (a) the role of the psychosocial domain in weight loss and weight management; (b) the guiding principles used to select constructs and measures for recommendation; (c) the initial high-priority constructs recommended for inclusion; (d) domain-specific issues for advancing the science; and (e) recommendations for future research.

Role of the Psychosocial Domain

In the ADOPT process-oriented model of treatment for weight loss (4), constructs within the psychosocial domain are hypothesized to serve at least one of three functions: predictor, mechanism (i.e., mediator), or moderator. A process-oriented approach, as operationalized by ADOPT, seeks to explicitly model and test the pathways through which these factors lead to successful treatment. As a predictor, psychosocial factors are associated with weight loss and weight maintenance, an association that is mediated through behavior or biology. In most cases, predictors will be constructs measured at baseline that predict treatment outcome, regardless of whether the construct changes due to the intervention. Psychosocial factors may also serve as a mechanism through which an intervention is effective. Interventions are often designed to target and change specific psychosocial inputs to, in turn, change the behavior that will have a direct effect on weight. For example, self-efficacy is a frequently studied target for treatment that has the potential to trigger a cascade of behaviors leading to weight loss: increasing self-efficacy increases healthy behaviors (e.g., diet, physical activity) that over time lead to successful weight loss (5). A distinction can be made between predictors and mechanisms: psychosocial predictors are constructs that predict treatment outcomes but do not necessarily change as a result of the intervention, whereas psychosocial mechanisms are changes in constructs that contribute to outcomes. It is important to note that a construct can be both a predictor at baseline and a mechanism through which changes in the construct mediate weight loss. For example, higher perceived stress at baseline may predict worse treatment outcomes (predictor), whereas declines in perceived stress in response to treatment may lead to better outcomes (mediator). Finally, factors within this domain may also serve as moderators of intervention effectiveness. Just as specific intervention strategies may be more effective for men than women (6), certain interventions may be more effective based on aspects of the individual's psychosocial functioning. Tailored interventions for physical activity based on an individual's motivation, for example, may be more effective for those who show enjoyment toward physical activity at baseline (7). Consistent with a precision medicine approach (8), identifying what type of intervention works best based on psychosocial functioning is critical for applying the most effective treatment approach. Theoretically, nearly all of the identified constructs in the psychosocial domain could operate as predictor, mechanism, and moderator. More extensive datasets generated through the ADOPT approach will help to better identify the specific functions the recommended psychosocial constructs are most likely to serve.

Guiding Principles

The psychosocial domain offers a rich range of constructs to explore individual differences in response to treatment. The inclusion of the same constructs/measures across interventions permits comparisons and will help identify constructs most important for effective interventions. Since we rarely have the benefit of direct comparisons between constructs, it is expected that this initial list will be refined by the scientific community over time, and these recommendations should not limit investigators to only these constructs/measures.

Construct selection

Constructs were selected based on their hypothesized relevance to weight loss/maintenance and the strength of the current empirical evidence. A construct would ideally be rooted in a theoretical model of processes that underlie weight loss/maintenance and be grounded in evidence of its utility from randomized controlled trials (RCTs) of weight-loss interventions. Constructs with evidence of predictive validity from either weight-loss/maintenance trials or trials that targeted key behaviors implicated in weight loss (e.g., diet, physical activity) were thus selected for inclusion. Some constructs within this domain, however, have a strong theoretical basis for weight loss/maintenance but have not yet been tested in an RCT (e.g., "wanting"/"liking"). Theoretically meaningful constructs with robust evidence from longitudinal studies of body weight and relevant behaviors were also prioritized.

Measure selection

We used another set of guiding principles to evaluate and select candidate measures. Reliability (e.g., test-retest, internal consistency) and validity of the measure were the primary considerations; after that, measures were selected to minimize participant and researcher burden. To reduce participant burden, we prioritized shorter scales that retained strong psychometric properties. Likewise, when possible, measures that are available publicly were chosen over measures under copyright or otherwise proprietary to help minimize cost to the investigator and maximize availability. We also considered ease of administration and scoring. For example, constructs measured with specific tasks may be preferable under some circumstances, but the expertise required to administer and score the task correctly makes it difficult to implement broadly in trials. As such, the self-report measure of a construct was chosen when there was a choice between a psychometrically-sound measure compared to a complex task that would challenge feasible implementation within a clinical trial. There is the expectation, however, that content experts (e.g., researchers knowledgeable about the construct and the scale used to measure it, such as individuals who have published on the construct) need to be consulted to aid with interpretation.

It is important to note that each of the ADOPT domains is not rigidly defined, and thus constructs could easily fall into more than one domain. Some eating-related behaviors, for example, include both engagement in the specific behavior and psychological antecedents that are a significant part of the behavior and thus could be placed in either the behavioral domain or the psychosocial domain. Constructs such as perceptions of satiety and hunger, for example, involve internal judgments (psychological) and actions based on those judgments (behavioral). Likewise, the underlying drivers of specific behaviors, such as binge eating, are often psychological in nature. These constructs were placed within the psychosocial domain

due to their psychological antecedents. There were also constructs that could be considered within the psychosocial domain that were placed within another domain. Most notably, social support and related social constructs (e.g., social norms) that capture aspects of the individual's social environment were included under the environmental domain. These constructs could have been included as part of the psychosocial domain but were placed in the environmental domain because they capture social aspects of the individual's surroundings. There are valid arguments as to why some constructs would be better placed in one domain or another. To reduce redundancy, however, constructs were only included once across the entire ADOPT framework. The complexity of where to place specific constructs highlights the need to consider all four domains within the ADOPT framework simultaneously.

High-Priority Constructs and Measures

Through a comprehensive review of the literature, expertise in aspects of psychosocial factors related to body weight, and the guiding principles described, we identified an initial list of high-priority constructs and their measures. See Table 1 for the recommended constructs and their associated recommended measures. The recommended constructs/measures are organized into four subdomains: (1) Affect, Stress, and Nonhomeostatic Eating, (2) Executive Function, (3) Motivation, and (4) Personality. Further information regarding each construct and its associated measure described below is available at the ADOPT Core Measures Workspace in the Grid-Enabled Measures (GEM) database (<https://www.gem-measures.org/workspaces/ADOPT>). GEM (www.gem-measures.org) is an electronic database of common measures in clinical research that is publically accessible and easy to use. Workspaces in GEM allow for input and discussion that help to build consensus around common measures.

Affect, stress, and nonhomeostatic eating

Nonhomeostatic eating, or eating beyond caloric need, includes common eating behaviors that contribute to weight gain. Chronic stress can increase nonhomeostatic eating and promote obesity and preferential deposition of visceral fat (9). Thus, it is theoretically important to include measures of distress and nonhomeostatic eating behavior (stress eating, binge eating, cravings, and reward-based eating). Further, stress can increase cravings and impair people's capabilities for adhering to interventions (10).

Affect and subjective stress. Perceived stress has been associated with nonhomeostatic eating behaviors in only a few cross-sectional studies. Given the theoretical importance of current perceived stress, it is recommended as part of ADOPT. The Perceived Stress Scale (11), a 4- or 10-item scale, measures feelings of stress over the last month.

State negative affect or acute stressful events have stronger links to nonhomeostatic eating than either perceived stress or trait affect. They tend to trigger overeating episodes in both animals and humans (12). State affect can be measured with the Positive and Negative Affect Schedule. Studies using ecological momentary assessment (EMA) (13) support the affect regulation model of binge

eating, which poses that negative affect precedes binges and, in turn, that bingeing tends to proximally reduce negative affect (14). Positive affect may trigger overeating episodes rather than binges (15) and needs further study. Daily measures are high in burden but offer promise for understanding individual differences in stress eating and identifying targets for intervention. For researchers able to measure daily (at the end of the day) or momentary (several times a day) responses, it may be helpful to include items for state negative affect (Positive and Negative Affect Schedule), cravings (see below), and episodes of overeating (see below) as intervention targets (e.g., see Boggiano et al. (16)).

Emotional eating. The Coping Subscale of the Palatable Eating Motives Scale is a 5-item subscale that measures intentionally using palatable food to cope with negative feelings (17). The trait form has been validated using EMA (16), and both versions are related equally to BMI and changes in emotional eating correlate with changes in BMI over time (18).

Binge eating. Binge eating disorder (BED) is recognized as a psychiatric disorder in the *Diagnostic and Statistical Manual of Mental Disorders, 5th Edition*. Criteria include eating an objectively large amount of food accompanied by loss of control (LOC) over eating at least weekly for 3 months. Many people with BED have obesity, and the prevalence of BED increases with increasing BMI and among those seeking weight-management treatment. The Questionnaire on Eating and Weight Patterns-5 (19), included in the PhenX toolkit (www.phenxtoolkit.org), has been adapted to reflect *Diagnostic and Statistical Manual of Mental Disorders, 5th Edition* criteria for BED and can be used to screen for its diagnosis. The item on binge eating frequency, which may reflect severity, can be used to see if change in this measure predicts treatment success. It is optional but strongly suggested that the items measuring subjective binge eating (feeling LOC without eating an objectively large amount) be used to learn if this also predicts poor treatment response. Further research is needed to determine whether a continuous measurement score of binge eating severity is predictive of outcomes for those who report binge or LOC eating, regardless of whether they meet criteria for BED.

Cravings and reward-based eating. There are many related aspects of nonhomeostatic eating, such as cravings, LOC, and lack of satiety. We recommend two measures: one for eating behaviors linked to reward drive and one for cravings. The Reward-Based Eating Scale (RED) is a 13-item scale that assesses preoccupation with food, LOC over eating, and lack of satiety with one reliable factor. Based on Item Response Theory, it includes items to provide greater coverage of the middle to low levels of reward-based eating commonly reported in obesity (20). The RED predicts weight cycling; weight gain over years (21); and, in one weight-loss trial, improvement in scores on the RED mediated weight (22). Food cravings, in the absence of hunger, are strong stimuli that can interfere with dietary intentions. The Trait Food Craving Questionnaire is a 15-item measure of behavioral, cognitive, and physical aspects of cravings for different types of food. High scores predict how much people crave and eat densely caloric snacks in daily life (23) and are associated with self-reported failures in dieting (24).

TABLE 1 Initial list of recommended constructs and measures from psychosocial domain subgroup of ADOPT working group

Construct	Measure	Logistics				Source of evidence
		Measure type	Number of items	Administration time (min)	Measurement schedule	
Affect, stress, and nonhomeostatic eating						
State affect	Positive and Negative Affect Schedule (state)	EMA, SA	20	< 5	B, MA	L
Perceived stress	Perceived Stress Scale	SA	4, 10	< 5	B, F	C
Emotional eating	Palatable Eating Motives: Coping Subscale	EMA, SA	4/full	< 5 / < 10	B, MA, F	L
Binge eating	Questionnaire on Eating and Weight Patterns	SA	11	< 5	B, F	C
Trait food craving	Trait Food Craving Questionnaire-Reduced	SA	15	< 5	B, F	C
Reward-related eating	Reward-Based Eating Drive	SA	13	< 5	B, F	RCT
Executive function						
Executive function	Behavior Rating Inventory of Executive Function-Adult	SA	75	< 30	B	C
Delay discounting	Kirby questionnaire	SA	27	< 10	B	C
Motivation						
Behavioral intention	Behavioral Intention Scale(s)	EMA, SA	Variable	Variable	B, MA	L
Self-efficacy	Self-Efficacy Scale(s)	EMA, SA	Variable	Variable	B, MA	L
Hedonic response to ("liking") and motivation ("wanting") for food	"Liking" and "Wanting" Visual Analog Scales	EMA, SA	2	< 5	B, MA	L
Hunger and satiety						
Hunger and satiety	Hunger and Satiety Visual Analog Scales	EMA, SA	2	< 5	B, MA	L
Personality						
Personality: Big Five factors	Mini-International Personality Item Pool (short form)/Big Five Inventory-2 (long form)	SA	20/60	< 5 / < 10	B, F	L

Evidence included both weight-related outcomes (e.g., weight loss) and behaviors related to weight loss (e.g., physical activity). B, baseline; C, cross-sectional; EMA, ecological momentary assessment; F, follow-up.; L, longitudinal; MA, multiple assessments across trial; RCT, randomized controlled trial; SA, self-administered.

Executive function

Executive function (EF) is considered a central concept in behavior control (25) and is critical for goal-directed behavior and the control of complex cognition (26). There is general agreement that there are three core EFs (27): inhibition, also called inhibitory control, including self-control (behavioral inhibition) and interference control (selective attention and cognitive inhibition); working memory; and cognitive flexibility (also called set shifting or mental flexibility). From these core functions, higher-order EFs are built, including reasoning, decision-making, problem solving, and planning (28). EF is necessary for successful weight loss, since it facilitates the ability to plan meals and physical activity, organize schedules, follow through on goals, inhibit desires, choose healthy foods despite the prevalence of easily accessible high-calorie foods, problem solve situations that could lead to overeating or lack of physical activity, and diligently self-monitor (29).

It is important to note that obesity could impact cognitive functioning, as comorbidities of obesity such as depression, hypertension, dyslipidemia, and type 2 diabetes are associated with cognitive impairment (30-32). Emerging animal and human research, however, suggests an independent contribution of obesity to cognitive impairment via a range of potential mechanisms, including impaired cerebral metabolism (33), elevated leptin (34), and inflammation and neuronal degradation (35).

Research suggests that EF deficits are found among individuals with overweight or obesity compared to those who are healthy weight (35,36). The evidence suggests that individuals with overweight or obesity, compared to those with healthy weight, have impairments in psychomotor performance and speed, verbal memory, set shifting, decision-making (including delay discounting), and inhibition (37-39). To date, there is no available evidence to support an independent relationship between obesity and working memory, learning, visual memory, and verbal fluency (36). It is important to note that studies have been inconsistent in delineating the effects of obesity versus obesity-related comorbidities, particularly depression and cardiovascular disorders, on cognition. Recent data also suggest that weight loss and/or exercise may improve memory, cognitive flexibility, and attention/EF (40).

There are numerous tasks that assess specific aspects of EF, and a comprehensive assessment of overall EF would require multiple tasks and take at least a few hours. For this reason, and the goal of maximizing reliability and validity while minimizing participant and researcher burden, and being sensitive to researcher expertise, we recommend the Behavior Rating Inventory of Executive Function-Adult Version (BRIEF-A). The BRIEF-A is a questionnaire composed of 75 items within 9 clinical scales that can relate to overeating, self-control, physical activity, and weight gain: Inhibit, Self-Monitor, Plan/Organize, Shift, Initiate, Task Monitor, Emotional Control, Working Memory, and Organization of Materials. Two broad indexes (Behavioral Regulation and Metacognition), an overall summary score, and three validity scales (Negativity, Inconsistency, and Infrequency) are also computed. The BRIEF-A has strong psychometric properties (41) and has been used in individuals with a range of conditions across a wide age range.

Delay discounting. Delay discounting, or temporal discounting, is a cognitive process that represents decision-making, specifically a

comparison of values between the immediate and delayed consumption of a determined commodity (42). Delay discounting depicts the tendency of people to give greater value to rewards as they move away from their temporal horizons and toward the present, with high delay discounting suggesting a preference for small, immediately available rewards as compared to larger rewards in the future (43). Although not typically included as a measure of EF per se, decision-making is considered a higher-order EF, and delay discounting represents a cognitive neuroscience perspective on decision-making in the context of reward.

Although delay discounting can be measured in a number of ways, the 27-item Kirby questionnaire is most commonly used. This questionnaire is easy to administer, meets psychometric standards for reliability and stability, and has 1-year test-retest correlations similar to tests designed to assess personality traits (44). Traditionally, delay discounting is measured using a discounting rate (k -parameter in a hyperbolic discounting function). However, recent studies suggest that a proportion of choices of the delayed rewards is relatively easy to calculate and highly correlated with the k -parameters (45). Reviews suggest that delay discounting could be a promising predictive measure of health behavior, as high discount rates for money are associated with several unhealthy behaviors and markers of health status (46). At this time, there is an emerging relation between delay discounting and weight loss (47); however, the data on a relation between delay discounting and BMI is mixed (48,49).

Motivation

Based on dual-processing theories (50), motivation can be divided into constructs that are a function of controlled processing (e.g., behavioral intention and self-efficacy) and constructs that are a function of automatic processing (e.g., “liking” and “wanting”) (51). Hunger and satiety are psychobiological motivational states that are also important determinants of eating behavior.

Behavioral intention and self-efficacy. Behavioral intention is how much someone wants, plans, or desires to perform a given behavior (52). Self-efficacy is the perceived capability to perform a given behavior (53). Behavioral intention and self-efficacy are a function of controlled cognitive processing regarding the consequences of a given behavior and the perceived facilitators and impediments to the behavior (52,53). We focus on behavioral intention because it is posited to be the most proximal determinant of behavior and thus a stronger predictor of behaviors that lead to weight loss than causal antecedents, such as attitudes, social norms, and perceived behavioral control (52). Self-efficacy is posited to impact behavior through its effects on outcome expectancies, social support, and perceived barriers (53). Emerging research, however, suggests that self-reported self-efficacy (i.e., what respondents say they “can do”) may reflect motivation to perform the target behavior rather than literal perceived capability and is thus likely to be—like behavioral intention—a proximal determinant of behavior (54).

Numerous scales have been developed to assess behavioral intentions (55) and self-efficacy (56) for weight loss or weight-loss-related behaviors. In choosing a measure of behavioral intention or self-efficacy, it is important that the measure be relevant to the particular behavior (e.g., exercise, eating fruits and vegetables), population (e.g., older adults, adults with overweight/obesity), and context (e.g., worksite, grocery shopping, rural, urban). Because of the

contextual nature of these measures, the evidence base for predictors of weight loss and weight-loss-related behaviors is best understood on the construct level (i.e., intention and self-efficacy) rather than specific measures of the construct. Reviews of behavioral intention and self-efficacy have shown that both constructs predict physical activity (57), dietary behaviors such as fruit and vegetable consumption (58), and weight loss (59).

“Liking” and “wanting”. The automatic processing aspects of motivation center around the concept of reward, which can be split into the neurobiological distinctions of “liking” and “wanting” (60). “Liking” refers to neurobiological underpinnings of the immediate pleasurable response to a stimulus, such as the taste of foods. “Wanting” refers to the neurobiological underpinnings of incentive salience—a specific kind of motivation mediated by the mesolimbic dopaminergic system and automatically triggered by presentation of a relevant behavioral cue. “Wanting” and “liking” are distinct phenomena, with “wanting” occurring before potential consumption of the reward and “liking” occurring during or after consumption. While previous “liking” of a stimulus generally leads to future “wanting” of that stimulus the next time it is cued, this relation is context-dependent such that there are instances in which “liking” does not lead to future “wanting” or “wanting” occurs without previous “liking” (60). Similar to the distinction between “liking” and “wanting,” “disliking” and “dread” are the negative poles of affective valence and incentive salience, respectively (60). Reference to these constructs always includes quotation marks to illustrate that these terms represent core neurobiological processes and thus may not map directly on to the broader concepts of liking and wanting (or disliking and dread).

The concepts of “liking” and “wanting” as components of reward are fundamental to understanding motivation for eating behavior and thus weight loss/maintenance (61) and can be measured with “Liking” and “Wanting” Visual Analog Scales. These visual analog scales—in which respondents are asked “Do you like [or want] this food?”—can be used to assess self-reported “liking” and “wanting” of foods associated with weight gain and/or targets of a planned weight-loss intervention, such as sugar-sweetened beverages, high-fat snack foods, or high-fat and high-sugar desserts. Higher ratings of “liking” and “wanting” for such foods has been observed in individuals with obesity (62). Additionally, the reward value of food is known to increase during short-term energy deficit (63), whereas changes in food “liking” and “wanting” have been shown to differentiate between those susceptible versus resistant to exercise-induced weight loss (64). Moreover, reductions in reward-driven eating have also been shown to account for some of the positive impacts of behavioral interventions on weight loss (22). Likewise, several studies have shown that assessment of “liking”/“disliking” is predictive of adherence to physical activity programs (65).

Hunger and satiety. Hunger is a subjective sensation that reflects a conscious motivation to eat; satiety is the suppression of hunger during the intermeal period that inhibits further eating (66). These processes are conceptualized through the satiety cascade, which provides a theoretical framework that maps the underlying biological mechanisms of appetite onto the psychological experience and behaviors that influence food intake (67). Subjective hunger and satiety are measured using horizontal visual analog scales anchored at each end using the extremes of a unipolar question relating to

subjective hunger, fullness, desire to eat, or prospective food consumption (68). Within-subject ratings are sensitive to experimental manipulation of food ingestion and display good test-retest reliability (69).

Measures of subjective hunger and satiety can be used to help explain eating behavior and susceptibility to weight loss or gain. An inability to recognize and respond to internal sensations of hunger (70) or a weakened satiety response to food consumption (71) are risk factors for overconsumption and weight gain. Hunger, which increases following weight loss and may remain elevated during weight-loss maintenance (72), is also predictive of successful long-term weight-loss maintenance (71).

Personality

Psychological dispositions are stable characteristics that describe a person’s typical ways of thinking, feeling, and behaving. The Five-Factor Model of personality (FFM) (73), also known as the Big Five, operationalizes these characteristics along five broad dimensions: Conscientiousness, Neuroticism, Extraversion, Openness, and Agreeableness.

Of the five traits, conscientiousness is associated most consistently with body weight and weight-related behaviors. Conscientiousness is defined as the general tendency to be organized, disciplined, and dutiful (74). Individuals who score high in conscientiousness are at less risk of weight gain and lower risk of obesity across adulthood (75,76). There is likewise a consistent association between conscientiousness and greater engagement in physical activity (77-79), consumption of a healthier diet (80-82), and healthier eating patterns (82). Of note, the associations between conscientiousness and BMI and behavior tend to be similar regardless of the measure used and tend to be similar across different demographic groups (76,78).

Neuroticism is the general tendency to experience negative emotions and greater vulnerability to stress (83). Neuroticism has been associated with greater weight gain in adulthood, particularly among women (75), but evidence is mixed (76). Neuroticism is, however, associated consistently with physical activity and eating patterns: Individuals higher in neuroticism are less physically active (77,78) and more likely to engage in disordered eating behavior (82,84). As such, although the evidence for a direct association between neuroticism and weight outcomes is somewhat equivocal, there is a clear association between neuroticism and behaviors conducive to obesity.

The other traits are more consistently associated with behaviors related to diet and physical activity than to body weight itself. There are very consistent associations, for example, between extraversion (the tendency to be outgoing and sociable) and more physical activity (77-79), less engagement in sedentary behavior (78), and consumption of a healthier diet (80-82). Openness (the tendency to be creative and unconventional) is likewise associated with more frequent physical activity (77,78) and, of the FFM traits, is the one associated most consistently and most strongly with consuming a healthy diet (80-82). There is not strong evidence that agreeableness (the tendency to be sympathetic and trusting) is related consistently with diet, physical activity, or weight outcomes (76,78,81), but it is strongly related to adherence in other behavioral domains (85). Thus, although not directly relevant to weight, it may be extremely

consequential for identifying individuals who are most (and least) likely to adhere to an intervention.

There are many reliable and valid measures of FFM personality traits. Common scales include the suite of Neuroticism-Extraversion-Openness (NEO) measures (e.g., NEO Personality Inventory 3, NEO Five-Factor Inventory (86)), the International Personality Item Pool (IPIP) (87), the Big Five Inventory (BFI) (88), the Midlife Development Inventory (89), and the Ten Item Personality Inventory (90). Each of these inventories is an excellent choice to measure personality traits. In keeping with our goal of maximizing reliability and validity while minimizing participant and researcher burden, we recommend the mini-IPIP as a measure of personality to include in adult weight-loss trials. The mini-IPIP is brief (it measures 5 traits with 20 items); has good psychometric properties (91); and is associated reliably with body weight (76), behaviors implicated in effective weight loss (78), and adherence (92). When time permits, we recommend the BFI-2. As with the mini-IPIP, the BFI-2 has excellent psychometric properties and predictive validity (88). The advantage of the BFI-2 is that it allows for the scoring of more specific aspects, or facets, of the broad traits. Facet-level analyses are able to help pinpoint exactly which aspects of the traits are associated with weight-loss outcomes and will thus help to both identify potential intervention targets and better understand the mechanisms through which personality traits lead to specific outcomes. The BFI-2 is a 60-item measure and thus should be used when the investigator and participants have more time to complete questionnaires.

Domain-Specific Considerations for Advancing the Science

We identified an initial list of high-priority constructs with a strong evidence base, but there is still much to be learned about how these constructs operate within trials, how they relate to each other, and how they relate to other key biological, behavioral, and environmental factors. The key next step is the consistent, coordinated measurement of these constructs (and those in the other domains) in trials for weight loss/maintenance. Such an approach will bring us closer to understanding individual and subgroup differences in response to treatment and to developing more targeted, and thus more effective, interventions. Addressing gaps in knowledge will provide a roadmap for future research that will help better identify how psychosocial factors contribute to weight loss/management and how to match interventions to the individual to increase the likelihood of success.

Evidence

Across the psychosocial domain, there is variability in the strength of evidence for specific constructs. Some constructs have substantial evidence from prospective longitudinal studies but lack evidence from RCTs. Conscientiousness, for example, is a consistent predictor of healthier weight trajectories across the lifespan (75,93), but the effect of this trait has not been tested in randomized trials. Constructs were recommended because of their theoretical relevance to weight and the strength of the evidence from observational studies. Still, their utility needs to be tested in the context of RCTs. The evidence from these RCTs will provide essential guidance for future

recommendations regarding high-priority constructs and measures in this domain.

Sociocultural factors

There are sociocultural factors that may contribute to the success or failure of interventions. The experience of interpersonal aggression, for example, has been associated with worse weight management. There is growing evidence that unfair treatment on the basis of body weight (94) and unfair treatment more generally (95) are associated with greater weight gain over time. Weight stigma in particular has been implicated in both weight gain and behaviors conducive to obesity and thus should be included in future work in clinical trials to determine its utility in the context of weight loss and maintenance. Such work will also be helpful in identifying the best measure of stigma to be recommended to be included in trials. In addition, other constructs within the psychosocial domain are present across a wide range of populations but vary in their association with weight-related outcomes by population (e.g., body image, attitudes toward weight (96)). Such constructs need to be tested further to determine their utility in weight-loss trials.

More generally, even when targeting the same construct or behavior, the intervention needs to take into account the sociocultural context of the individual being treated. Interventions that are culturally sensitive and tailored to a population tend to be more effective than interventions designed for other populations or treatment as usual (97). Several of the recommended constructs have been associated with weight-related factors in multiple populations (e.g., personality, stress), but the effectiveness of interventions based on these constructs may vary across populations and needs to be explicitly tested.

Health disparities

There are well-documented disparities in obesity prevalence across demographic groups, including racial and ethnic minority, rural, and low-income populations (98). And yet, these groups are under-represented in clinical trial research (99), especially weight-loss trials (100). RCTs tend to focus on white, urban, middle-class individuals. It is critical to include more diverse populations, especially populations that have the greatest disparities in obesity, in clinical trials. Weight-loss trials are also needed that test interventions tailored to hypothesized population-specific needs. These efforts will allow for a more robust scientific base explaining the predictors, mediators, and moderators unique to groups at the highest risk for obesity and help identify targets to improve treatment approaches for these populations. Although ADOPT encourages the use of consistent measures across studies, testing whether constructs important for weight loss in one group generalize to others is also critical.

Feasibility of measures/technology

As technology progresses, there will be more opportunities to integrate sophisticated assessments into weight-loss trials. This issue is usually of concern in domains such as biology and behavior but also applies to psychosocial constructs. For example, as the ability to more easily incorporate EMA into trials increases, there are more opportunities to measure constructs in real time and intervene on

aspects of psychological functioning that fluctuate regularly throughout the day (e.g., affect, stress). Such advances in technology may also facilitate study into interactions across domains, such as how psychosocial functioning and biology reciprocally influence each other. An example is the potential for use of sensor cueing used to direct individuals to alternative strategies rather than eating to self-soothe in response to stress.

Process/mechanisms

In the development and delivery of obesity treatments, knowledge of psychosocial processes is important for understanding which intervention strategies modify the behavioral and biological factors that have a direct effect on weight loss and weight maintenance. These constructs also represent the thoughts and feelings people have in response to changes in their weight; responses that, in turn, affect people's ability to engage in behaviors that enable them to maintain a weight-reduced state. Because of the dynamic and interactive nature of these constructs across domains, the ADOPT initiative encourages investigators to include low-burden measures from each psychosocial subdomain when possible. This approach will facilitate research into the dynamic relation between domains to identify how to best intervene for weight loss. The ADOPT effort is a step toward an integrated process model that explicitly recognizes and encourages research into interconnections to advance the science of effective intervention for weight loss.

Recommendations for Research

Measurement schedule

How frequently a construct needs to be measured in a trial depends on how the construct is hypothesized to operate. Constructs hypothesized to be predictors or moderators should be measured at the beginning of the trial. Constructs that are hypothesized to be mediators of intervention effectiveness need to be measured at multiple points throughout the trial. Further, some constructs are hypothesized to be predictors but may also change over the course of the trial because of either the intervention or through the process of weight loss, and that change may be the mechanism that explains the effectiveness of the intervention. Even constructs hypothesized to be predictors or moderators may turn out to be mediators if change in the construct contributes to better outcomes. Thus, the constructs often need to be measured multiple times over the course of the trial. Table 1 shows the suggested measurement schedule for the recommended measures. These recommendations are based on the current state of the evidence; future work needs to better specify these schedules and ground them in a better understanding of how these constructs change over time.

Promising constructs

In addition to our initial set of recommended measures, it is imperative to continue to identify how psychosocial factors contribute to effective weight loss/management. Our recommendations are meant to help bring consistency to weight-loss trials to promote comparisons and integration across trials. These recommendations should not, however, limit investigations into additional psychosocial factors implicated in weight loss.

Conclusions

The purpose of this paper was to describe the psychosocial domain and its importance to the ADOPT project, to define the constructs and measures recommended for inclusion in trials, and to place this domain within the broader weight loss/maintenance process. The complexity of weight loss underscores the importance of addressing the multiple factors that contribute independently and in combination to healthy weight regulation. The inclusion of similar measures across different weight-loss trials will help bring coherence to the literature and facilitate comparisons and integration of data across trials. An added benefit is that by including measures from the four domains in a single trial, hypotheses about connections and interactions across domains can be tested to identify how different domains function together to improve or hinder intervention effectiveness. Such evidence is critical to advance the design and delivery of more effective treatments for weight loss and weight-loss maintenance. **O**

© 2018 The Obesity Society

References

- Santos I, Sniehotta FF, Marques MM, Carraça EV, Teixeira PJ. Prevalence of personal weight control attempts in adults: a systematic review and meta-analysis. *Obes Rev* 2017;18:32-50.
- Dombrowski SU, Knittle K, Avenell A, Araújo-Soares V, Sniehotta FF. Long term maintenance of weight loss with non-surgical interventions in obese adults: systematic review and meta-analyses of randomised controlled trials. *BMJ* 2014;348:g2646. doi: 10.1136/bmj.g2646
- MacLean PS, Wing RR, Davidson T, et al. NIH working group report: innovative research to improve maintenance of weight loss. *Obesity (Silver Spring)* 2015;23:7-15.
- MacLean PS, Rothman AJ, Nicastro HL, et al. The Accumulating Data to Optimally Predict obesity Treatment (ADOPT) Core Measures Project: rationale and approach. *Obesity (Silver Spring)* 2018;26(suppl 2):S6-S15.
- Warziski MT, Sereika SM, Styn MA, Music E, Burke LE. Changes in self-efficacy and dietary adherence: the impact on weight loss in the PREFER study. *J Behav Med* 2008;31:81-92.
- Williams RL, Wood LG, Collins CE, Callister R. Effectiveness of weight loss interventions—is there a difference between men and women: a systematic review. *Obes Rev* 2015;16:171-186.
- Williams D, Papandonatos G, Napolitano M, Lewis B, Whiteley J, Marcus B. Perceived enjoyment moderates the efficacy of an individually tailored physical activity intervention. *J Sport Exercise Psychol* 2006;28:300-309.
- Collins FS, Varmus H. A new initiative on precision medicine. *N Engl J Med* 2015;372:793-795.
- Mason A, Epel E. Stress and obesity. In: Brownell K, Walsh T, eds. *Eating Disorders and Obesity: A Comprehensive Handbook*. New York, NY: Guilford Press; 2017.
- Sinha R, Jastreboff AM. Stress as a common risk factor for obesity and addiction. *Biol Psychiatry* 2013;73:827-835.
- Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav* 1983;24:385-396.
- Razzoli M, Pearson C, Crow S, Bartolomucci A. Stress, overeating, and obesity: insights from human studies and preclinical models. *Neurosci Biobehav Rev* 2017;76:154-162.
- Engel SG, Crosby RD, Thomas G, et al. Ecological momentary assessment in eating disorder and obesity research: a review of the recent literature. *Curr Psychiatry Rep* 2016;18:37. doi: 10.1007/s11920-016-0672-7
- Berg KC, Crosby RD, Cao L, Crow SJ, Engel SG, Wonderlich SA, Peterson CB. Negative affect prior to and following overeating-only, loss of control eating-only, and binge eating episodes in obese adults. *Int J Eat Disord* 2015;48:641-653.
- Sultson H, Kukk K, Akkermann K. Positive and negative emotional eating have different associations with overeating and binge eating: construction and validation of the Positive-Negative Emotional Eating Scale. *Appetite* 2017;116:423-430.
- Boggiano MM, Wenger LE, Turan B, et al. Real-time sampling of reasons for hedonic food consumption: further validation of the Palatable Eating Motives Scale. *Front Psychol* 2015;6:744. doi: 10.3389/fpsyg.2015.00744
- Burgess EE, Turan B, Lokken KL, Morse A, Boggiano MM. Profiling motives behind hedonic eating. Preliminary validation of the Palatable Eating Motives Scale. *Appetite* 2014;72:66-72.

18. Boggiano MM, Wenger LE, Turan B, Tatum MM, Morgan PR, Sylvester MD. Eating tasty food to cope. Longitudinal association with BMI. *Appetite* 2015;87:365-370.
19. Yanovski SZ, Marcus MD, Wadden TA, Walsh BT. The Questionnaire on Eating and Weight Patterns-5: an updated screening instrument for binge eating disorder. *Int J Eat Disord* 2015;48:259-261.
20. Mason AE, Vainik U, Acree M, Tomiyama AJ, Dagher A, Epel ES, Hecht FM. Improving assessment of the spectrum of reward-related eating: the RED-13. *Front Psychol* 2017;8:795. doi: 10.3389/fpsyg.2017.00795
21. Epel ES, Tomiyama AJ, Mason AE, et al. The reward-based eating drive scale: a self-report index of reward-based eating. *PLoS One* 2014;9:e101350. doi: 10.1371/journal.pone.0101350
22. Mason AE, Epel ES, Aschbacher K, et al. Reduced reward-driven eating accounts for the impact of a mindfulness-based diet and exercise intervention on weight loss: data from the SHINE randomized controlled trial. *Appetite* 2016;100:86-93.
23. Richard A, Meule A, Reichenberger J, Bleichert J. Food cravings in everyday life: an EMA study on snack-related thoughts, cravings, and consumption. *Appetite* 2017;113:215-223.
24. Meule A, Lutz A, Vögele C, Kübler A. Food cravings discriminate differentially between successful and unsuccessful dieters and non-dieters. Validation of the Food Cravings Questionnaires in German. *Appetite* 2012;58:88-97.
25. Stuss DT, Knight RT. Introduction. In: Stuss DT, Knight RT, eds. *Principles of Frontal Lobe Function*. New York: Oxford University Press; 2002:1-7.
26. Banich MT. Executive function: the search for an integrated account. *Curr Dir Psychol Sci* 2009;18:89-94.
27. Miyake A, Friedman NP, Emerson MJ, Witzki AH, Howerter A, Wager TD. The unity and diversity of executive functions and their contributions to complex "frontal lobe" tasks: a latent variable analysis. *Cognitive Psychol* 2000;41:49-100.
28. Collins A, Koechlin E. Reasoning, learning, and creativity: frontal lobe function and human decision-making. *PLoS Biol* 2012;10:e1001293. doi: 10.1371/journal.pbio.1001293
29. Foster GD, Makris AP, Bailer BA. Behavioral treatment of obesity. *Am J Clin Nutr* 2005;82:230S-235S.
30. Panza F, D'Introno A, Colacicco AM, et al. Lipid metabolism in cognitive decline and dementia. *Brain Res Rev* 2006;51:275-292.
31. Whitmer RA, Sidney S, Selby J, Johnston SC, Yaffe K. Midlife cardiovascular risk factors and risk of dementia in late life. *Neurology* 2005;64:277-281.
32. Areosa SA, Grimley EV. Effect of the treatment of type II diabetes mellitus on the development of cognitive impairment and dementia. *Cochrane Database Syst Rev* 2002:CD003804. doi: 10.1002/14651858.CD003804
33. Volkow ND, Wang GJ, Telang F, et al. Inverse association between BMI and prefrontal metabolic activity in healthy adults. *Obesity (Silver Spring)* 2009;17:60-65.
34. Gunstad J, Spitznagel MB, Keary TA, et al. Serum leptin levels are associated with cognitive function in older adults. *Brain Res* 2008;1230:233-236.
35. Smith E, Hay P, Campbell L, Trollor JN. A review of the association between obesity and cognitive function across the lifespan: implications for novel approaches to prevention and treatment. *Obes Rev* 2011;12:740-755.
36. Prickett C, Brennan L, Stolwyk R. Examining the relationship between obesity and cognitive function: a systematic literature review. *Obes Res Clin Pract* 2015;9:93-113.
37. Weller RE, Cook EW, Avsar KB, Cox JE. Obese women show greater delay discounting than healthy-weight women. *Appetite* 2008;51:563-569.
38. Davis C, Pate K, Curtis C, Reid C. Immediate pleasures and future consequences. A neuropsychological study of binge eating and obesity. *Appetite* 2010;54:208-213.
39. Gunstad J, Paul RH, Cohen RA, Tate DF, Spitznagel MB, Gordon E. Elevated body mass index is associated with executive dysfunction in otherwise healthy adults. *Compr Psychiatry* 2007;48:57-61.
40. Siervo M, Arnold R, Wells JC, Tagliabue A, Colantuoni A, Albanese E, Brayne C, Stephan BC. Intentional weight loss in overweight and obese individuals and cognitive function: a systematic review and meta-analysis. *Obes Rev* 2011;12:968-983.
41. Roth RM, Isquith PK, Gioia GA. *BRIEF-A: Behavior Rating Inventory of Executive Function-Adult Version*. Lutz, FL: Psychological Assessment Resources; 2005.
42. Lowenstein G. Anticipation and the valuation of delayed consumption. *Econom J* 1987;97:666-684.
43. Tesch AD, Sanfey AG. Models and methods in delay discounting. *Ann N Y Acad Sci* 2008;1128:90-94.
44. Kirby KN. One-year temporal stability of delay-discount rates. *Psychon Bull Rev* 2009;16:457-462.
45. Myerson J, Baumann AA, Green L. Discounting of delayed rewards: (A)theoretical interpretation of the Kirby questionnaire. *Behav Processes* 2014;107:99-105.
46. Story GW, Vlaev I, Seymour B, Darzi A, Dolan RJ. Does temporal discounting explain unhealthy behavior? A systematic review and reinforcement learning perspective. *Front Behav Neurosci* 2014;8:76. doi: 10.3389/fnbeh.2014.00076
47. Sze YY, Slaven EM, Bickel WK, Epstein LH. Delay discounting and utility for money or weight loss. *Obes Sci Pract* 2017;3:69-74.
48. Barlow P, Reeves A, McKee M, Galea G, Stuckler D. Unhealthy diets, obesity and time discounting: a systematic literature review and network analysis. *Obes Rev* 2016;17:810-819.
49. Lim SL, Bruce AS. Can't wait to lose weight? Characterizing temporal discounting parameters for weight-loss. *Appetite* 2015;85:8-13.
50. Hofmann W, Friese M, Wiers RW. Impulsive versus reflective influences on health behavior: A theoretical framework and empirical review. *Health Psychol Rev* 2008;2:111-137.
51. Williams DM, Evans DR. Current emotion research in health behavior science. *Emotion Rev* 2014;6:277-287.
52. Fishbein M. A reasoned action approach to health promotion. *Med Decis Making* 2008;28:834-844.
53. Bandura A. Health promotion by social cognitive means. *Health Ed Behav* 2004;31:143-164.
54. Williams DM, Rhodes RE. The confounded self-efficacy construct: conceptual analysis and recommendations for future research. *Health Psychol Rev* 2016;10:113-128.
55. Cruwys T, Platow MJ, Rieger E, Byrne DG. The development and validation of the Dieting Intentions Scale (DIS). *Psychol Assess* 2013;25:264-278.
56. Wilson KE, Harden SM, Almeida FA, You W, Hill JL, Goessl C, Estabrooks PA. Brief self-efficacy scales for use in weight-loss trials: Preliminary evidence of validity. *Psychol Assess* 2016;28:1255-1264.
57. Rhodes RE, Quinlan A. Predictors of physical activity change among adults using observational designs. *Sports Med* 2015;45:423-441.
58. Reyes BF, Warner LM, Knoll N, Montenegro E, Schwarzer R. Synergistic effects of social support and self-efficacy on dietary motivation predicting fruit and vegetable intake. *Appetite* 2015;87:330-335.
59. Delahanty LM, Peyrot M, Shrader PJ, et al. Pretreatment, psychological, and behavioral predictors of weight outcomes among lifestyle intervention participants in the Diabetes Prevention Program (DPP). *Diabetes Care* 2013;36:34-40.
60. Berridge KC, Robinson TE. Parsing reward. *Trends Neurosci*. 2003;26:507-513.
61. Finlayson G, King N, Blundell JE. Liking vs. wanting food: importance for human appetite control and weight regulation. *Neurosci Biobehav Rev* 2007;31:987-1002.
62. Berthoud HR, Zheng H, Shin AC. Food reward in the obese and after weight loss induced by calorie restriction and bariatric surgery. *Ann N Y Acad Sci* 2012;1264:36-48.
63. Epstein LH, Truesdale R, Wojcik A, Paluch RA, Raynor HA. Effects of deprivation on hedonics and reinforcing value of food. *Physiol Behav* 2003;78:221-227.
64. Finlayson G, Caudwell P, Gibbons C, Hopkins M, King N, Blundell J. Low fat loss response after medium-term supervised exercise in obese is associated with exercise-induced increase in food reward. *J Obes* 2011;2011:615624. doi: 10.1155/2011/615624
65. Rhodes RE, Kates A. Can the affective response to exercise predict future motives and physical activity behavior? A systematic review of published evidence. *Ann Behav Med* 2015;49:715-731.
66. Blundell J, De Graaf C, Hulshof T, et al. Appetite control: methodological aspects of the evaluation of foods. *Obes Rev* 2010;11:251-270.
67. Blundell J. Pharmacological approaches to appetite suppression. *Trends Pharmacol Sci* 1991;12:147-157.
68. Rogers PJ, Blundell JE. Effect of anorexic drugs on food intake and the micro-structure of eating in human subjects. *Psychopharmacology* 1979;66:159-165.
69. Stubbs R, Hughes D, Johnstone A, et al. The use of visual analogue scales to assess motivation to eat in human subjects: a review of their reliability and validity with an evaluation of new hand-held computerized systems for temporal tracking of appetite ratings. *Brit J Nut* 2000;84:405-415.
70. Barkeling B, King NA, Näslund E, Blundell JE. Characterization of obese individuals who claim to detect no relationship between their eating pattern and sensations of hunger or fullness. *Int J Obes (Lond)* 2007;31:435-439.
71. Drapeau V, King N, Hetherington M, Doucet E, Blundell J, Tremblay A. Appetite sensations and satiety quotient: predictors of energy intake and weight loss. *Appetite* 2007;48:159-166.
72. Sumithran P, Prendergast LA, Delbridge E, et al. Long-term persistence of hormonal adaptations to weight loss. *N Engl J Med* 2011;365:1597-1604.
73. McCrae RR, Costa PT. The Five-Factor Theory of personality. In: John OP, Robins RW, Pervin LA, eds. *Handbook of Personality: Theory and Research*. 3rd ed. New York, NY: Guilford Press; 2008.
74. Roberts BW, Lejuez C, Krueger RF, Richards JM, Hill PL. What is conscientiousness and how can it be assessed? *Dev Psychol* 2014;50:1315-1330.
75. Sutin AR, Ferrucci L, Zonderman AB, Terracciano A. Personality and obesity across the adult life span. *J Pers Soc Psychol* 2011;101:579-592.
76. Jokela M, Hintsanen M, Hakulinen C, et al. Association of personality with the development and persistence of obesity: a meta-analysis based on individual-participant data. *Obes Rev* 2013;14:315-323.

77. Wilson KE, Dishman RK. Personality and physical activity: a systematic review and meta-analysis. *Pers Ind Diff* 2015;27:230-242.
78. Sutin AR, Stephan Y, Luchetti M, Artese A, Oshio A, Terracciano A. The five factor model of personality and physical inactivity: a meta-analysis of 16 samples. *J Res Pers* 2016;63:22-28.
79. Rhodes RE, Smith NEI. Personality correlates of physical activity: a review and meta-analysis. *Brit J Sport Med* 2006;40:958-965.
80. Möttus R, McNeill G, Jia X, Craig LC, Starr JM, Deary IJ. The associations between personality, diet and body mass index in older people. *Health Psychol* 2013;32:353-360.
81. Möttus R, Realo A, Allik J, Deary IJ, Esko T, Metspalu A. Personality traits and eating habits in a large sample of Estonians. *Health Psychol* 2012;31:806-814.
82. Sutin AR, Terracciano A. Personality traits and body mass index: modifiers and mechanisms. *Psychol Health* 2016;31:259-275.
83. Costa PT, Jr., McCrae RR. *Revised NEO Personality Inventory (NEO-PI-R) and the NEO Five-Factor Inventory (NEO-FFI) Professional Manual*. Odessa, FL: Psychological Assessment Resources; 1992.
84. Elfhag K, Morey LC. Personality traits and eating behavior in the obese: poor self-control in emotional and external eating but personality assets in restrained eating. *Eat Behav* 2008;9:285-293.
85. Axelsson M, Brink E, Lundgren J, Lötvalld J. The influence of personality traits on reported adherence to medication in individuals with chronic disease: an epidemiological study in West Sweden. *PLoS One* 2011;6:e18241. doi: 10.1371/journal.pone.0018241
86. McCrae RR, Costa PT. *NEO Inventories for the NEO Personality Inventory-3, NEO Five-Factor Inventory-3, and NEO Personality Inventory-Revised Professional Manual*. Lutz, FL: Psychological Assessment Resources; 2010.
87. Goldberg LR, Johnson JA, Eber HW, et al. The international personality item pool and the future of public domain personality measures. *J Res Pers* 2006;40:84-96.
88. Soto CJ, John OP. The next Big Five Inventory (BFI-2): developing and assessing a hierarchical model with 15 facets to enhance bandwidth, fidelity, and predictive power. *J Pers Soc Psychol* 2017;113:117-143.
89. Lachman ME, Weaver SL. *Midlife Development Inventory (MIDI) Personality Scales: Scale Construction and Scoring. Technical report*. Waltham, MA: Brandeis University; 1997.
90. Gosling SD, Rentfrow PJ, Swann WB. A very brief measure of the big-five personality domains. *J Res Pers* 2003;37:504-528.
91. Donnellan MB, Oswald FL, Baird BM, Lucas RE. The mini-IPIP scales: tiny-yet-effective measures of the Big Five factors of personality. *Psychol Assess* 2006;18:192-203.
92. Moran AM, Everhart DE, Davis CE, Wuensch KL, Lee DO, Demaree HA. Personality correlates of adherence with continuous positive airway pressure (CPAP). *Sleep Breath* 2011;15:687-694.
93. Sutin AR, Kerr JA, Terracciano A. Temperament and body weight from ages 4 to 15 years. *Int J Obes (Lond)* 2017;41:1056-1061.
94. Sutin AR, Terracciano A. Perceived weight discrimination and obesity. *PLoS One* 2013;8:e70048. doi: 10.1371/journal.pone.0070048
95. Hunte HE. Association between perceived interpersonal everyday discrimination and waist circumference over a 9-year period in the Midlife Development in the United States cohort study. *Am J Epidemiol* 2011;173:1232-1239.
96. Robinson SA, Webb JB, Butler-Ajibade PT. Body image and modifiable weight control behaviors among black females: a review of the literature. *Obesity (Silver Spring)* 2012;20:241-252.
97. Barrera M, Castro FG, Strycker LA, Toobert DJ. Cultural adaptations of behavioral health interventions: a progress report. *J Consult Clin Psychol* 2013;81:196-205.
98. Wang Y, Beydoun MA. The obesity epidemic in the United States—gender, age, socioeconomic, racial/ethnic, and geographic characteristics: a systematic review and meta-regression analysis. *Epidemiol Rev* 2007;29:6-28.
99. Oh SS, Galanter J, Thakur N, et al. Diversity in clinical and biomedical research: a promise yet to be fulfilled. *PLoS Med* 2015;12:e1001918. doi: 10.1371/journal.pmed.1001918
100. Kumanyika SK, Whitt-Glover MC, Haire-Joshu D. What works for obesity prevention and treatment in black Americans? Research directions. *Obes Rev* 2014;15(suppl 4):204-212.